

Erythema

Erythema	Disorder
annulare marginatum	rheumatic fever by β -hemolyzing A streptococci (ASL, anti-hyaluronidase, anti-DNase)
chronicum migrans	borreliae (IgG, IgM ab)
contagiosum	parvo virus B19 (IgG, IgM ab), PCR
exudativum multiforme	Herpes virus (IgG, IgM ab), PCR
infectiosum acutum	parvo virus B19 (IgG, IgM ab), PCR
nodosum	rheumatic fever by β -hemolyzing A streptococci (see above), M. Boeck (ACE), coccidiomycosis (ab), yersiniae, (O3, O9 ab, bacteriol.), leprosy (histol.), TBC (PCR, bacteriol.), colitis ulcerosa, histoplasmosis, Beçet syndrome, lymphogranuloma venereum (Chlamydia trachomatis PCR), contraceptives (e.g. ethinylestradiol), M. Crohn, for), drug induced (e.g. penicillin, ampicillin, sulfonamide, iodide bromide)
subitum	parvo virus B19, herpes virus type 6 (Ab; PCR), echo virus, coxsackie virus, Morbus Still
toxicum neonatorum	eosinophilic granulocytes increased in vesicles, Giemsa staining recommended (DD: staphylo-dermia)

For complete list of laboratory test offered at Freiburg Medical Laboratory, please visit <http://www.fml-dubai.com/parameter-listings/>