



Freiburg Medical Laboratory Middle East (L.L.C.)

P.O. Box: 3068, Dubai - UAE, Tel: 04 396 2227, Fax: 04 396 2228

E-mail: info@fml-dubai.com, Website: www.fml-dubai.com

Physician:

Dr. M. Jaksch
Freiburg Medical Lab

Laboratory Report Online Version

Report Date: 15.11.2018

Patient Name: NBS normal

Gender: Female

Date of Birth: 13.11.2018

Nationality:

Your ID:

Test Request Code: 1810

Sample ID:

Patient IDNo: 380518

Sampling Date / Time: 15.11.2018 / 00:00

Receipt Date / Time: 15.11.2018 / 18:24

Remarks:

Insurance:

Analysis	Result	Flag	Units	Reference Range
----------	--------	------	-------	-----------------

Exclusion of Metabolic Diseases (Filter card)

Week of gestation 39

Newborn screening* See text

No sign of the following disorders:

Amino Acid related disorders such as: Phenylketonuria (PKU),

Hyperphenylalaninaemia, Maple syrup disease etc.

Fatty acid oxidation disorders and other disorders recognized in the acylcarnitine spectrum: e.g. MCAD, LCAD, VLCAD

Other screened disorders: e.g. Hypothyreosis, AGS, Biotinidase deficiency,

Galactosaemia, G6PDH Deficiency, Cystic Fibrosis, Hemoglobinopathies

(HbS, HbC, HbD, HbE, HbH/HbBarts, HbO, HbLepore, HbG, beta-Thalassaemia)

Methods: MS-MS, FIA, Photometry, Capillary Electrophoresis

Please note that the best time of collecting blood for NBS is 36-72 hours after birth.

Amino Acids (PKU etc.)*	normal	quantitative	normal
Acylcarnitines (Fatty acids)*	normal	quantitative	normal
TSH (Hypothyreodism)*	normal	quantitative	normal
17a-OH-Progesterone (AGS)*	normal	quantitative	normal
Galactose (Galactosemia)*	normal	quantitative	normal
Biotinidase (Biot.deficiency)*	normal	quantitative	normal
Glucose-6-PDH*	normal	quantitative	normal
Immunoreactive Trypsin*	normal	quantitative	normal
Hemoglobin Screening*	normal	quantitative	normal

Note:

Our reference values are adjusted to age and gender.

Daily internal Quality Control within the required range

(according to ISO 15189).

External Quality Control available on request.

^ non-accredited parameter

*This parameter is affected by Biotin intake of >5 mg (RDI = 0.03mg)

* This investigation has been performed in a collaborating accredited laboratory (Germany).

Techn. Validation by
Med. Technologist
(Supervisor of
the Department)

Dr. Nehmat ElBanna
Specialist
Clinical Pathology (U/S)
(DHA-P-0084548)

PD Dr. med. habil. M. Jaksch
Associate Professor
Medical Director
(DHA-LS-240710)

This report has been printed through lmed-Online-Reporting-System and therefore does not carry a signature.