

Oxidative capacity

General:

Oxidative stress: Oxidative stress means the discrepancy between formation and resorption of free radicals (reactive oxygen species (ROS)). Exogeneous sources of ROS are smoking, alcohol consumption, air pollution, chemical toxins, heavy metals, ozone, UV rays. Endogeneous sources of ROS are inflammations, immune defense, stress, chronic diseases and excessive physical efforts. Environmental poisons, medication etc. lead to an increased formation of free radicals in the body (oxidative capacity).

Free radicals: Free radicals react directly with cellular structures such as lipids, proteins and nucleic acids. The reaction with nonsaturated fatty acids in lipid membranes is of high significance due to increased lipid peroxidation. This can cause cell damage. The body usually reacts with repair mechanisms (antioxidants) which can be tested and differentiated in the an-tioxidative capacity.

Antioxidants: Antioxidants are natural or synthetic substances which protect cells and food from damage by action of oxygen radicals (free radicals). Especially vitamin C and E, the provitamin beta-carotin as well as certain trace elements, e.g. selenium, coenzyme Q10 are among the natural anti-oxidants contained in food.

Synthetic antioxidants (e.g. propylgallate, gallate, butyl-hydroxy-toluol orthophosphoric acid, phosphate and others) delay the oxidation of fatty acids. In potato products and other vegetables they prevent discolorations caused by the air oxygen. Antioxidants in food must be declared mentioning the sub-stance and the E number.

Enzyme Redox systems	Material
Malondialdehyde (MDA)	EDTA plasma frozen
Superoxide dismutase (SOD)	Heparin blood
Glutathione reduced (GSHR)	EDTA
Glutathione total (GSHT)	EDTA

Vitamins and trace elements	Material
Vitamin E (Tocopherol, VITE)	2 ml of serum (light protected)
Vitamin C (ascorbic acid, VITC)	1 ml heparin plasma (light protected)
β -Carotin (CARO)	2 ml of serum (light protected)
Zinc (ZN)	5 ml of serum
Coenzyme Q10 (COQ10)	2 ml of serum
Selenium (SE)	5 ml heparin blood
Ceruloplasmin (COER)	1 ml of serum

The following tests are available:

- **Complete oxidative capacity**

Indication: Determination of the oxidative load

Material: 1 ml EDTA-plasma, frozen

TAT: 7-10 days*

Method: photometry

Units: $\mu\text{mol/l}$

Ref.- range: normal oxidative load: < 200 $\mu\text{mol/l}$
 moderate oxidative load: 200 - 350 $\mu\text{mol/l}$
 strong oxidative load: > 350 $\mu\text{mol/l}$

- **Total antioxidative capacity**

General:

This screening tests the antioxidative balance and includes antioxidative protection factors.

Indication: high infect sensitivity, neurological disorders, gastrointestinal disorders etc., optimization of the therapy with antioxidants or diet etc.

Material: 1 ml EDTA-plasma, frozen

TAT: 7-10 days*

Method: photometry

Units: $\mu\text{mol/l}$

Ref.- range: low antioxidative capacity: < 280 $\mu\text{mol/l}$

medium antioxidative capacity: 280 - 320 $\mu\text{mol/l}$

high antioxidative capacity: > 320 $\mu\text{mol/l}$

• **Other parameters**

Enzyme Redox systems

	Material
Malondialdehyde	Serum
Superoxide dismutase	EDTA-blood
Glutathione, reduced	CPDA-blood
Glutathione, total	CPDA-blood

Vitamins and trace elements

	Material
Vitamin E (Tocopherol)	Serum, light-protected
Vitamin C (Ascorbic acid)	Serum, frozen
Beta-Carotene	Serum, light-protected
Zinc	Serum
Coenzyme Q10	Serum
Selenium	Serum or LH-blood (trace-free tubes)
Ceruloplasmin	Serum

For complete list of laboratory test offered at Freiburg Medical Laboratory, please visit
<http://www.fml-dubai.com/parameter-listings/>